



Responses of terrestrial ecosystems and carbon budgets to current and future environmental variability

Citation

Medvigy, D., S. C. Wofsy, J. W. Munger, and P. R. Moorcroft. 2010. "Responses of Terrestrial Ecosystems and Carbon Budgets to Current and Future Environmental Variability." Proceedings of the National Academy of Sciences 107, no. 18: 8275–8280.

Published Version

doi:10.1073/pnas.0912032107

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:13041311>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

PNAS

^aDepartment of Organismic and Evolutionary Biology and ^bSchool of Engineering and Applied Sciences and Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA 02138

We assess the significance of high-frequency variability of environmental parameters (sunlight, precipitation, temperature) for the structure and function of terrestrial ecosystems under current and future climate. We examine the influence of hourly, daily, and monthly variance using the Ecosystem Demography model version 2 in conjunction with the long-term record of carbon fluxes measured at Harvard Forest. We find that fluctuations of sunlight and precipitation are strongly and nonlinearly coupled to ecosystem function, with effects that accumulate through annual and decadal timescales. Increasing variability in sunlight and precipitation leads to lower rates of carbon sequestration and favors broad-leaved deciduous trees over conifers. Temperature variability has only minor impacts by comparison. We also find that projected changes in sunlight and precipitation variability have important implications for carbon storage and ecosystem structure and composition. Based on Intergovernmental Panel on Climate Change model estimates for changes in high-frequency meteorological variability over the next 100 years, we expect that terrestrial ecosystems will be affected by changes in variability almost as much as by changes in mean climate. We conclude that terrestrial ecosystems are highly sensitive to high-frequency meteorological variability, and that accurate knowledge of the statistics of this variability is essential for realistic predictions of ecosystem structure and functioning.

Changes in climate expected in the next century (1) are likely to have important consequences for the structure, composition, and functioning of terrestrial ecosystems (2–4). Mean climate (5, 6) and climate variability are projected to shift in concert (7); for example, warmer climate may be associated with greater variability in the northeastern United States (8), increasing variability of rainfall intensity during the South Asian monsoon (9), and increasing frequency of heat waves (10).

In this paper, we assess the influence of meteorological variability on ecosystem function and development for typical forest ecosystems in the northeastern United States, using the Ecosystem Demography model version 2 (ED2) biosphere model (24, 25) and nearly 10 years of eddy-flux measurements. We assess how strongly this system responds to high-frequency variability of sunlight, temperature, and precipitation, and identify the most important statistical measures and the underlying mechanisms for response. ED2 realistically simulates the physiological functioning, growth, death, and recruitment of individual plants and of the whole forest ecosystem, including regional structure and vegetation dynamics, for timescales from hours to decades (*Materials and Methods*) (24, 25). We find large, systematic differences in ecosystem functioning and the resulting structure and composition of the forest when ED2 is driven by observed hourly meteorological forcing as compared with meteorological drivers that have the same mean values but fail to reproduce high-frequency temporal variation, including sophisticated products generally considered suitable for ecosystem-climate studies.

ED2 was initialized using observed stand composition and carbon stocks at Harvard Forest, and driven with 10 years of hourly meteorological data from the site (S_{full}), monthly averaged hourly mean values of the same data (S_{mm} ; Fig. 1), and other datasets (see below). Removal of the high-frequency variability enhanced decadal net ecosystem productivity (NEP) by 50%, from 4.6 tons of carbon per hectare per year ($\text{tC ha}^{-1} \text{y}^{-1}$) to $3.1 \text{ tC ha}^{-1} \text{y}^{-1}$. Both gross primary productivity (GPP) and total ecosystem respiration (R_{tot}) were artificially elevated (Fig. 1), but the effect was much larger for GPP (16%; $15.4 \text{ tC ha}^{-1} \text{y}^{-1}$ versus $13.3 \text{ tC ha}^{-1} \text{y}^{-1}$) than for R_{tot} (6%). A similar nonlinear response was found for the whole northeastern region in a 100-yr integration (Fig. 24): Mean NEP was artificially enhanced by more than 50% ($1.0 \text{ tC ha}^{-1} \text{y}^{-1}$ versus $0.63 \text{ tC ha}^{-1} \text{y}^{-1}$) (SI Text and Fig. S1).

ECOLOGY

Based on simulations driven by AOGCM output (Table 2), we propose here that climate change will impact terrestrial ecosystems in part through changes in high-frequency meteorological variability. The strength of this effect may be comparable to that arising from changes in mean meteorological drivers. However, uncertainties in climate models limit our confidence in quantitative estimates of changes in future variability and associated carbon fluxes.

Our simulations were all carried out with a single ecosystem model that, like all models, is an incomplete and imperfect representation of reality. Additional empirical tests are needed to further evaluate and develop the mechanisms proposed here. Specifically, multiyear records from eddy-flux towers at other sites, including conifer-dominated sites, could be used to test whether conifers do, in fact, exhibit a larger response to variability than deciduous trees. And given the strong effect of canopy interception on ecosystem functioning, additional measurements of throughfall effects would be very useful for model evaluation.

We conclude that studies of climate-ecosystem interactions require careful representation of meteorological forcing, including their high-frequency statistical variances. This requirement becomes more stringent as biosphere/land-surface models become more realistic and as dynamic changes in vegetation come into focus. The mechanisms identified here whereby meteorological variances influence ecosystem responses on hourly-to-monthly timescales are general, applying to forests throughout the globe, and hence the requirements for models and model drivers are, likewise, broadly applicable. Changes in high-frequency as well as low-frequency climate variability will have important consequences for the composition, structure, and functioning of terrestrial ecosystems.

Materials and Methods

Harvard Forest Simulations. The ED2 model is described in *SI Text*. Physical and biogeochemical soil properties needed to initialize the Harvard Forest simulations were obtained from field measurements. The initial forest structure and composition were obtained from the forest inventory measurements within the flux-tower footprint following ref. 25.

The meteorological drivers required by ED2 are solar radiation, long-wave radiation, temperature, humidity, precipitation, wind speed, and pressure. For the Harvard Forest simulations, these were specified from the EMS eddy-flux tower meteorological observations, with any gaps filled by measurements from a nearby weather station. The simulation S_{full} used the hourly mean values of all meteorological drivers. S_{mm} was driven by the monthly-hourly mean of the meteorological forcing used in S_{full} . The GISS, CNRM, GFDL, ECMWF, and NCEP output had temporal resolutions ranging from 1,800 (GISS) to 10,800 (CNRM, GFDL) to 21,600 (ECMWF, NCEP) seconds. Instantaneous values of all meteorological drivers except solar radiation were

generated from a simple linear interpolation in time. Instantaneous values of solar radiation were obtained by weighting the radiation values in the original datasets by the cosine of the solar zenith angle. The ISCCP simulation was identical to S_{full} , except that it used 3-hourly ISCCP radiation linearly interpolated in time. All meteorological fields were rescaled to match the observed monthly-hourly mean values, yielding forcing datasets differing only in their higher-order statistics.

Regional Simulations. All regional simulations were done on the $0.5^\circ \times 0.5^\circ$ grid shown in Fig. 2A. Soil textural class was assigned at the level of the grid cell using the $1^\circ \times 1^\circ$ resolution US Department of Agriculture global soil database because higher-resolution data were unavailable for Quebec. Forest inventory data (43, 44) were used to initialize the forest composition. Horizontal heterogeneity was captured by defining each inventory plot as a separate patch within each grid cell, whereas the vertical heterogeneity within each plot was captured by defining each tree as a cohort with species assignment to the appropriate plant functional type.

Meteorological drivers for the regional simulations were obtained from the ECMWF ERA-40 reanalysis (28), which had a temporal resolution of 6 h. These 6-hourly values were disaggregated into hourly values using the same procedure used in the Harvard Forest simulations (see above). The regional S_{mm} was driven by monthly mean diurnal cycles of the meteorological forcing used in the regional S_{full} .

Following ref. 25, the vegetation phenology was prescribed from moderate-resolution imaging spectroradiometer-derived estimates of leaf onset and offset dates averaged between 2001 and 2004 (45). Spatial patterns of forest harvesting were derived from forest inventory data and were applied as a disturbance forcing to the model using the methodology of ref. 17. The period June 1982–June 2082 was simulated, but the first 5 simulated years (1982–1986) were used only to equilibrate the soil carbon pools and to allow for the establishment of grasses (which are not accounted for in the forest inventories) in recently harvested patches.

ACKNOWLEDGMENTS. We thank Mark Friedl and Tan Bin for their assistance and generosity in providing the moderate-resolution imaging spectroradiometer (MODIS)-derived leaf onset and offset dates used in the phenology model. We acknowledge the modeling groups, the Program for Climate Model Diagnosis and Intercomparison (PCMDI), and the World Climate Research Programme (WCRP)'s Working Group on Coupled Modelling (WGCM) for their roles in making available the WCRP CMIP3 multimodel dataset. Support of this dataset is provided by the Office of Science, US Department of Energy. S.C.W., J.W.M., and P.R.M. gratefully acknowledge funding from National Science Foundation BE/CBC Program Grant ATM-0221850 and BE Program Grant ATM-0450307. S.C.W., J.W.M., and P.R.M. also acknowledge funding from the Harvard Forest Long Term Ecological Research site, and from the Office of Science, Biological and Environmental Research Program (BER) of the US Department of Energy, through the northeast regional center of the National Institute for Global Environmental Change (NIGEC), and through the northeastern region of the National Institute for Climatic Change Research (NICCR) under Cooperative Agreement DE-FC02-03ER63613. Measurements at the Harvard Forest site are supported by the BER through the NICCR (DE-FC02-06ER64157) and the Terrestrial Carbon Program (DE-FG02-07ER64358).

- Meehl GA, et al. (2007) *Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, Global Climate Projections. Climate Change 2007: The Physical Science Basis*, eds Solomon S, et al. (Cambridge Univ Press, Cambridge, UK), pp 747–845.
- Cox PM, Betts RA, Jones CD, Spall SA, Totterdell IJ (2000) Acceleration of global warming due to carbon-cycle feedbacks in a coupled climate model. *Nature* 408:184–187.
- Melillo JM, et al. (2002) Soil warming and carbon-cycle feedbacks to the climate system. *Science* 298:2173–2176.
- Hanson PJ, et al. (2005) Importance of changing CO₂, temperature, precipitation, and ozone on carbon and water cycles of an upland-oak forest: Incorporating experimental results into model simulations. *Glob Change Biol* 11:1402–1423.
- Smith TM, Reynolds RW (2005) A global merged land-air-sea surface temperature reconstruction based on historical observations (1880–1997). *J Clim* 18:2021–2036.
- Vose RS, Wuertz D, Peterson TC, Jones PD (2005) An intercomparison of trends in surface air temperature analyses at the global, hemispheric, and grid-box scale. *Geophys Res Lett* 32:L18718.
- Easterling DR, et al. (2000) Climate extremes: Observations, modeling, impacts. *Science* 289:2068–2074.
- Wettstein JJ, Mearns LO (2002) The influence of the North Atlantic-Arctic Oscillation on the mean, variance, and extremes of temperature in the northeastern United States and Canada. *J Clim* 15:3586–3600.
- Goswami BN, Venugopal V, Sengupta D, Madhusoodanan MS, Xavier PK (2006) Increasing trend of extreme rain events over India in a warming environment. *Science* 314:1442–1445.
- Schar C, et al. (2004) The role of increasing temperature variability in European summer heatwaves. *Nature* 427:332–336.
- Marani M, Grossi G, Wallace M, Napolitano F, Entekhabi D (1997) Forcing, intermittency, and land surface hydrological partitioning. *Water Resour Res* 33:167–175.
- Margulis SA, Entekhabi D (2001) Temporal disaggregation of satellite-derived monthly precipitation estimates and the resulting propagation of error in partitioning of water at the land surface. *Hydrol Earth Syst Sci* 5:27–38.
- Ruel JJ, Ayres MP (1999) Jensen's inequality predicts effects of environmental variation. *Trends Ecol Evol* 14:361–366.
- McGuire AD, et al. (2001) Carbon balance of the terrestrial biosphere in the twentieth century: Analyses of CO₂, climate and land use effects with four process-based ecosystem models. *Global Biogeochem Cycles* 15:183–206.
- Cao M, et al. (2003) Response of terrestrial carbon uptake to climate interannual variability in China. *Glob Change Biol* 9:536–546.
- Tian H, et al. (1998) Effect of interannual climate variability on carbon storage in Amazonian ecosystems. *Nature* 396:664–667.
- Albani MA, Medvigy D, Hurtt GC, Moorcroft PR (2006) The contributions of land-use change, CO₂ fertilization, and climate variability to the eastern US carbon sink. *Glob Change Biol* 12:2370–2390.
- Nemani RR, et al. (2000) Climate-driven increases in global terrestrial net primary production from 1982 to 1999. *Science* 300:1560–1563.
- Berthelot M, Friedlingstein P, Ciais P, Dufresne J-L, Monfray P (2005) How uncertainties in future climate change predictions translate into future terrestrial carbon fluxes. *Glob Change Biol* 11:959–970.

20. Ito A (2005) Climate-related uncertainties in projections of the twenty-first century terrestrial carbon budget: Off-line model experiments using IPCC greenhouse-gas scenarios and AOGCM climate projections. *Clim Dyn* 24:435–448.
21. Gu L, et al. (2003) Response of a deciduous forest to the Mount Pinatubo eruption: Enhanced photosynthesis. *Science* 299:2035–2038.
22. Lee X, Wu H-J, Sigler J, Oishi C, Siccamo T (2004) Rapid and transient response of soil respiration to rain. *Glob Change Biol* 10:1017–1026.
23. Ensminger I, et al. (2004) Intermittent low temperatures constrain spring recovery of photosynthesis in boreal Scots pine forests. *Glob Change Biol* 10:995–1008.
24. Medvigy DM (2006) The state of the regional carbon cycle: Results from a coupled constrained ecosystem-atmosphere model. PhD thesis (Harvard University).
25. Medvigy D, Wofsy SC, Munger JW, Moorcroft PR (2009) Mechanistic scaling of ecosystem function and dynamics in space and time: Ecosystem Demography model version 2. *J Geophys Res* 114:G01002.
26. Sinclair TR, Murphy CE, Knoerr KR (1976) Development and evaluation of simplified models for simulating canopy photosynthesis and transpiration. *J Appl Ecol* 13:813–829.
27. Leuning R, Kelliher FM, de Pury DGG, Schulze E-D (1995) Leaf nitrogen, photosynthesis, conductance and transpiration: Scaling from leaves to canopies. *Plant Cell Environ* 18:1183–1200.
28. Uppala SM, et al. (2005) The ERA-40 re-analysis. *Q J R Meteorol Soc* 131:2961–3012.
29. Kalnay E, et al. (1996) The NCEP/NCAR 40-year reanalysis project. *Bull Am Met Soc* 77: 437–471.
30. Delworth TL, Knutson TR (2000) Simulation of early 20th century global warming. *Science* 287:2246–2250.
31. Schmidt GA, et al. (2006) Present day atmospheric simulations using GISS ModelE: Comparison to in-situ, satellite and reanalysis data. *J Clim* 19:153–192.
32. Salas-Méila D, et al. (2005) *Description and Validation of the CNRM-CM3 Global Coupled Model* (CNRM, Toulouse), Note de Centre du CNRM No 103.
33. Zhang Y, Rossow WB, Lacis AA, Oinas V, Mishchenko MI (2004) Calculation of radiative fluxes from the surface to top of the atmosphere based on ISCCP and other global data sets: Refinements of the radiative transfer model and the input data. *J Geophys Res* 109:D19105.
34. Pitman AJ, Henderson-Sellers A, Yang ZL (1990) Sensitivity of regional climates to localised precipitation in global models. *Nature* 364:734–747.
35. Wang D, Wang G, Anagnostou EN (2009) Impact of sub-grid variability of precipitation and canopy water storage on hydrological processes in a coupled land-atmosphere model. *Clim Dyn* 32:649–662.
36. Rödenbeck C, Houweling S, Gloor M, Heimann M (2003) CO₂ flux history 1982–2001 inferred from atmospheric data using a global inversion of atmospheric transport. *Atmos Chem Phys* 3:1919–1964.
37. Baker DF, et al. (2006) TransCom 3 inversion intercomparison: Impact of transport model errors on the interannual variability of regional CO₂ fluxes. *Global Biogeochem Cycles* 20:GB1002.
38. Peylin P, et al. (2005) Multiple constraints on regional CO₂ flux variations over land and oceans. *Global Biogeochem Cycles* 19:GB1011.
39. Kaminski T, Knorr W, Rayner PJ, Heimann M (2002) Assimilating atmospheric data into a terrestrial biosphere model: A case study of the seasonal cycle. *Global Biogeochem Cycles* 16:1066.
40. Scholze M, Kaminski T, Rayner P, Knorr W, Giering R (2007) Propagating uncertainty through prognostic carbon cycle data assimilation system simulations. *J Geophys Res* 112:D17305.
41. Denning AS, Fung IY, Randall D (1995) Latitudinal gradient of atmospheric CO₂ due to seasonal exchange with land biota. *Nature* 376:240–243.
42. Dupigny-Giroux L-A, Blackwell CF, Bristow S, Olson GM (2003) Vegetation response to ice disturbance and consecutive moisture extremes. *Int J Remote Sens* 24:2105–2129.
43. Frayer WE, Furnival GM (1999) Forest survey sampling designs: A history. *J For* 97: 4–10.
44. Penner M, Power K, Muirhead C, Tellier R, Wang Y (1997) *Canada's Forest Biomass Resources: Deriving Estimates from Canada's Forest Inventory* (Canadian Forest Service, Victoria, BC), Information Report No BC-X-370.
45. Zhang X, Friedl MA, Schaaf CB (2006) Global vegetation phenology from moderate resolution imaging spectroradiometer (MODIS): Evaluation of global patterns and comparison with in situ measurements. *J Geophys Res* 111:G04017.